Carbohydrate Research, 232 (1992) C13–C15
Elsevier Science Publishers B.V., Amsterdam

Contents

Conformational analysis of levanbiose by molecular mechanics
J. Liu (New Orleans, LA, USA) and A.L. Waterhouse (Davis, CA, USA) 1

Rearrangement of unsaturated 2,4-*O*-benzylidenehexitol derivatives into *C*-glycosylbenzene derivatives
J. Kuszmann, B. Podányi and G. Jerkovich (Budapest, Hungary) 17

The synthesis and characterisation of 2-*O*-(6-*O*-L-*glycero*-α,β-D-*manno*-heptopyranosyl-α-D-glucopyranosyl)- α,β-D-glucopyranose
S.A. Nepogodev, Z. Pakulski, A. Zamojski (Warszawa, Poland), O. Holst and H. Brade (Borstel, Germany) 33

Glucosylenamines as glycosyl acceptors: synthesis of gentiobiosylenamines
J. Fuentes Mota, J.F.-B. Guzmán, J.M. García Fernández, W. Moreda, C. Ortiz, M.A. Pradera, I. Robina and C. Welsh (Sevilla, Spain) 47

Synthesis of the antibiotic cortalcerone from D-glucose using pyranose 2-oxidase and a novel fungal enzyme, aldos-2-ulose dehydratase
K. Koths, R. Halenbeck and M. Moreland (Emeryville, CA, USA) 59

Studies of the interaction of the maltose-binding protein of *Escherichia coli*, a closed-groove binder, with 4,6-*O*-ethylidenemalto-oligosaccharides (dp 2–5) and its regioselective labelling with 3-azibutyl 1-thio-α-(6-^{3}H)maltoside
J. Lehmann, E. Schiltz and J. Steck (Freiburg, Germany) 77

Synthesis of 4′-deoxy-4′-fluorokanamycin A and B
Y. Takahashi, S. Tsuneda, T. Tsuchiya, Y. Koyama and S. Umezawa (Kawasaki, Japan) .. 89

Synthesis of a phenyl β-avobioside derivative of the disaccharide component of avoparcins
X. Dong, F. Tillequin, C. Monneret (Paris, France), G. Horvath and F. Sztaricskai (Debrecen, Hungary) 107

Free radical-mediated addition of peracetylated 1-bromo-β-D-glucopyranosyl chloride to acrylonitrile
J.-P. Praly, Z.E. Kharraf and G. Descotes (Villeurbanne, France) 117

The reaction of some dicarbonyl sugars with aminoguanidine
J. Hirsch (Columbia, MO, USA), E. Petrakova (Bethesda, MD, USA) and M.S. Feather (Columbia, MO, USA) 125

Immunochemical characterization of polyclonal and monoclonal *Streptococcus* group A antibodies by chemically defined glycoconjugates and synthetic oligosaccharides
K.B. Reimer (Burnaby, BC, Canada), M.A.J. Gidney, D.R. Bundle (Ottawa, ON, Canada) and B.M. Pinto (Burnaby, BC, Canada) 131

Structure of the polysaccharide O-antigen of *Salmonella riogrande* O:40 (group R) related to blood group A activity
M.B. Perry and L.L. MacLean (Ottawa, ON, Canada) 143

Formation of oligosaccharides from lactitol by *Aspergillus oryzae* β-D-galactosidase
S. Yanahira, T. Suguri, T. Yakabe, Y. Ikeuchi, G. Hanagata and E. Deya (Kawagoe, Japan) 151

Notes

Binding constants for complexes of α-cyclodextrin with L-phenylalanine and some related substrates
S. Chokchainarong, O.R. Fennema and K.A. Connors (Madison, WI, USA) 161

Synthesis 1-thioglycosides from 1-sulfenates
I. Fokt and W. Szeja (Gliwice, Poland) 169

Facile synthesis of 6-deoxy-L-*arabino*-hexulose (L-rhamnulose) 1-phosphate via regioselective phosphorylation of the unprotected sugar
P. Zhou, G. Lajoie, J.F. Honek, P.T.C. Nambiar and O.P. Ward (Waterloo, ON, Canada) 177

Synthesis of 2-amino-1,2-dideoxy-D-galactitol hydrochloride and 5-amino-1,4-anhydro-5,6-dideoxy-L-galactitol hydrochloride
K. Adelhorst and G.M. Whitesides (Cambridge, MA, USA) 183

Preliminary communication

Stereospecific 1,2-*cis*-glycosylation: a modified thiocyanate method
N.K. Kochetkov, E.M. Klimov, N.N. Malysheva and A.V. Demchenko (Moscow, Russia) . C1

Papers

Crystal structure of lactitol (4-O-β-D-galactopyranosyl-D-glucitol) trihydrate
J. Kivikoski, I. Pitkänen (Jyväskylä, Finland) and J. Nurmi (Kantvik, Finland) 189

The crystal structure and physicochemical properties of L-ascorbic acid 2-glucoside
T. Mandai, M. Yoneyama, S. Sakai, N. Muto and I. Yamamoto (Okayama, Japan) 197

Studies of the molecular recognition of synthetic methyl β-lactoside analogues by *Ricinus communis* agglutinin
A. Rivera-Sagredo, J. Jiménez-Barbero, M. Martín-Lomas, D. Solís and T. Díaz-Mauriño (Madrid, Spain) 207

Investigation of the active site of *Escherichia coli* β-D-galactosidase by photoaffinity labelling
C.-S. Kuhn, J. Lehmann (Freiburg i. Br., FRG), G. Jung and S. Stevanović (Tübingen, FRG) 227

Oxazolidines from sugars: conformationally restricted derivatives of muramic acid
J.M. Vega-Pérez, J.L. Espartero, F.J. Ruiz and F. Alcudia (Sevilla, Spain) 235

Some 3,5-acetal derivatives of D-glucitol
A.J. Al-Kadir, N. Baggett and J.M. Webber (Birmingham, UK) 249

Redox glycosidation: the use of Nozaki–Takai methylenylation in a highly stereoselective synthesis of sucrose
A.G.M. Barrett, L.M. Melcher (Fort Collins, CO, USA) and B.C.B. Bezuidenhoudt (Evanston, IL, USA) 259

Synthesis of 5-deoxy-5-fluoro and 5-deoxy-5,5-difluoro derivatives of kanamycin B and its analogs. Study on structure–toxicity relationships
T. Shitara, Y. Kobayashi, T. Tsuchiya and S. Umezawa (Kawasaki, Japan) 273

Chemical reactions induced by ultrasound and γ-rays in aqueous solutions of L-ascorbic acid
G. Portenlänger and H. Heusinger (Garching, FRG) 291

Apple-fruit xyloglucans: a comparative study of enzyme digests of whole cell walls and of alkali-extracted xyloglucans
C.M.G.C. Renard (Nantes, France), J.A. Lomax and J.J. Boon (Amsterdam, Netherlands) 303

Notes

Crystal and molecular structure of ethyl 6,7,8-tri-*O*-acetyl-5,9-anhydro-2,3,4-trideoxy-2-diphenylmethyleneamino- 4-nitro-D-*threo*-D-*altro*-nononate
P. Köll (Oldenburg, Germany), J. Kopf, D. Abeln (Hamburg, Germany), J. BeMiller and L. Petruš (West Lafayette, IN, USA) 321

Conformational analysis of the trisaccharide components of the repeating units of the capsular polysaccharides of *Streptococcus pneumoniae* types 19F and 19A
P. Ciuffreda, D. Colombo, F. Ronchetti (Milano, Italy) and L. Toma (Pavia, Italy) 327

Synthesis of methyl 3-amino-deoxy-D-alluronate
A. Gomtsyan, I. Savelyeva, S. Belyakov and I. Kalvinsh (Riga, Latvia) 341

Steric effect of a bulky 6-substituent in the I^+-promoted glycosylation with pent-4-enyl and thioethyl glycosides
S. Houdier and P.J.A. Vottero (Grenoble, France) 349

Synthesis and properties of 1,1,3,3-tetramethyl-2-(2,3,4,6-tetra-*O*-acetyl-α-D-glucopyranosyl)uronium triflate
K. Bock, J.F.-B. Guzmán (Valby, Denmark) and S. Refn (Lyngby, Denmark) 353

An improved preparation of 2,3,5-tri-*O*-acyl-β-D-ribofuranosyl azides by the Lewis acid-catalysed reaction of β-D-ribofuranosyl acetates and trimethylsilyl azide: an example of concomitant formation of the α anomer by trimethylsilyl triflate catalysis
A. Štimac (Novo mesto, Slovenia) and J. Kobe (Ljubljana, Slovenia) 359

Mechanism of the adsorption of pancreatic alpha-amylase onto starch crystallites
V.M. Leloup, P. Colonna (Nantes, France) and G. Marchis-Mouren (Marseille, France) . 367

O-(3-Butenyl), a stable blocking group removable by ozonolysis
C. Bayle, J. Defaye (Grenoble, France), D. Horton (Columbus, OH, USA), J. Lehmann and M. Scheuring (Freiburg i. Br., FRG) 375

Author index C7

Subject index C9

Contents (Vol. 232) C13